

COVID-19 VACCINATION CARD

Information about COVID-19 Vaccination

Depending on the COVID-19 vaccine you have received, it consists of two or three doses, and additional doses may be recommended. These must be completed to achieve the best possible protection, and for the protection to be as long-lasting as possible.

The vaccine has been assessed to be safe for use. However, just like other vaccines, you may experience some side effects such as headache, body aches, tiredness and soreness at the injection site, or fever. These usually get better after 1-3 days and may be a sign that your immune system is making a protective response against COVID-19.



MINISTRY OF HEALTH
SINGAPORE

COVID-19 VACCINATION CARD

Please bring this card for your next appointment

PERSONAL PARTICULARS

Name: _____

NRIC/FIN.: _____

DOB.: _____

YOUR NEXT VACCINATION IS DUE IN _____ DAYS

Appointment Date	Time	Place

COVID-19 VACCINATION RECORD

Vaccine Dose #	Date of Vaccination	Vaccine Brand	Place of Vaccination
		<input type="checkbox"/> Pfizer-BioNTech / Comirnaty (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Moderna / Spikevax (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Sinovac-CoronaVac <input type="checkbox"/> Novavax / Nuvaxovid	
		<input type="checkbox"/> Pfizer-BioNTech / Comirnaty (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Moderna / Spikevax (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Sinovac-CoronaVac <input type="checkbox"/> Novavax / Nuvaxovid	
		<input type="checkbox"/> Pfizer-BioNTech / Comirnaty (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Moderna / Spikevax (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Sinovac-CoronaVac <input type="checkbox"/> Novavax / Nuvaxovid	
		<input type="checkbox"/> Pfizer-BioNTech / Comirnaty (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Moderna / Spikevax (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Sinovac-CoronaVac <input type="checkbox"/> Novavax / Nuvaxovid	
		<input type="checkbox"/> Pfizer-BioNTech / Comirnaty (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Moderna / Spikevax (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Sinovac-CoronaVac <input type="checkbox"/> Novavax / Nuvaxovid	
		<input type="checkbox"/> Pfizer-BioNTech / Comirnaty (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Moderna / Spikevax (Original / Bivalent / XBB.1.5)* <input type="checkbox"/> Sinovac-CoronaVac <input type="checkbox"/> Novavax / Nuvaxovid	

*Specify version of vaccine administered

AFTER VACCINATION ADVICE

Possible Side Effects	How to Manage
Pain, redness, swelling at the injection site	Those with fever are advised to self-isolate at home until the fever subsides.
Fever, chills	
Headache, muscle pain, joint pain	Paracetamol 1 to 2 tablets every 6 hours for adults or dosed according to the child's weight as needed
Tiredness	Rest
Lymph node swelling at neck or arms	Usually gets better by itself in a week or so

See a doctor if:

- the side effects persist or get worse
- the fever persists for more than 48 hours

You can also report any side effects that you experience to HSA via: go.gov.sg/srf

This vaccine may cause a severe allergic reaction in very rare instances. If you experience a severe allergic reaction (**difficulty breathing, swelling of your face, throat, eyes or lips, a fast heartbeat, dizziness or weakness, rash all over your body**), please seek immediate medical attention by calling 995 or going to the nearest A&E. The doctor will also decide if you should receive the second dose of the vaccine.

If you experience chest pain, difficulty breathing or abnormal heartbeats, please also seek medical attention immediately.